




United States
Department of Agriculture
Forest Service, Idaho Falls, Idaho

March 2003

Poisonous Plants of South-East Idaho



The United States Department of Agriculture (USDA) prohibits discrimination in its programs on the basis of race, color, national origin, sex, religion, age, disability, political beliefs, and marital or familial status. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means of communication of program information (Braille, large print, audiotape, etc.) should contact the USDA Office of Communications at (202) 720-2791. To file a complaint, write the Secretary of Agriculture, U.S. Department of Agriculture, Washington, DC 20250, or call (202)720-7327 (voice) or (202)720-1127(TDD). USDA is an equal employment opportunity employer.

By: Wayne R. Owen, Boise National Forest
Alexis Collins, Environmental Education Coordinator
Revised for Southeast Idaho by:
Rose Lehman, Forest Botanist
Jared Mattson, Student, Idaho State University

Poison ivy	12
Spurge.....	10
Steer's head.....	9
Tansy	14
Tansy ragwort.....	13
Water Hemlock.....	7

Plants Especially Poisonous to Animals

Buttercup	12
Cherry	11
Cocklebur.....	16
Corn lily	15
Death camas	16
Dogbane.....	6
Klamath weed.....	10
Larkspur.....	9
Locoweed.....	7
Lupine.....	11
Milkweed	6
Poison hemlock.....	8
Spurge.....	10
Tansy ragwort.....	13
Water hemlock.....	7

Poisonous Plants of South-East Idaho

Every year, many adults and children become seriously ill by consuming poisonous plants. Sometimes this poisoning is accidental, and sometimes poisoning is the result of the misidentification or misuse of medicinal herbs. Our beloved animal companions (dogs, horses, llamas, etc.) may also fall victim to poisonous plants.

As a public service, the Caribou-Targhee National Forest has prepared this brief field guide to the poisonous plants commonly found in Southeast Idaho. We hope this information enhances your recreation experience. Most of the flowers and plants in the forest are harmless, so please don't think the woods are full of danger. Take the time to be informed and enjoy nature in a safe and informed way.

Although many of these plants are poisonous, many are attractive native wildflowers and are used by knowledgeable people for medicinal purposes.

For more information on poisonous plants, contact your public library, your local National Forest botanist, or one of these internet websites:

<http://www.fs.fed.us/r1/kootenai/resources/plants/botany/index.shtml>

http://www.vth.colostate.edu/poisonous_plants/report/search.cfm

<http://www.ansci.cornell.edu/plants/alphalist.html>

<http://plants.usda.gov>

<http://sis.agr.ca/brd/poisonpl/>

<http://www.pprl.usu.edu/>

<http://educ.ahsl.arizona.edu/poison.html>


If you are poisoned; call the local Poison Control Center at:

1-800-860-0620

Ferns and Their Allies


Horsetails and Scouring Rushes

(*Equisetum spp.*) Horsetails are common in moist, coarse-textured soils. The most common species in forested areas has “leafy” stems, but lowland varieties are typically leafless. Although the rough texture is enough to keep most people from eating them, young children and horses can eat enough to cause a stomach ache. American Indians and early settlers used these to scrub cooking utensils.



Braken Fern (*Pteridium aquilinum*)

This large fern can be found in forested areas and in dry upland openings around Idaho. The fronds and fiddleheads (emerging fronds) are poisonous unless thoroughly cooked. The fronds can be toxic to horses or llamas if consumed in quantity over a period of a week or more.



Index

Plants to Avoid Touching


Cocklebur (leaves)	16
Klamath weed (oily droplets on stem)	10
Milkweed (sap)	6
Poison ivy	12
Spurge	10
Stinging nettle	15

Plants Especially Poisonous to People

Baneberry	5
Braken fern	4
Buttercup	12
Corn lily	15
Death Camas	16
Elderberry	13
Horsetail	4
Hound's tongue	8
Klamath weed	10
Larkspur	9
Locoweed	7
Lupine	11
Monkshood	5
Nightshade	14
Poison hemlock	8


Cocklebur (*Xanthium strumarium*)

Cocklebur is a coarse weedy annual herb introduced to America from Europe. It is most often found along river banks and lake shores. No human poisonings from cocklebur are known but seeds and seedlings are toxic and potentially fatal to animals. People with sensitive skin may develop a rash from handling the plant. Cocklebur is sometimes confused with burdock (*Arctium spp.*) which is a larger plant having more spherical burs. Burdock may also cause dermatitis in sensitive individuals but it is not poisonous. Burdock root can be eaten as a raw vegetable.



Death Camas (*Zygadenus spp.*)

The grass-like leaves of death camas can be confused with wild onions (though death camas lacks the distinctive onion odor) and blue camas (death camas flowers are creamy white). This plant is common in grasslands and shrub lands throughout our area. The entire plant is highly toxic and fatal to both humans and animals. Poisoning symptoms include profuse salivation, burning lips, mouth numbness, thirst, stomach pain, vomiting, diarrhea, confusion, slow irregular heartbeat, low blood pressure and low temperature, difficulty breathing, coma and death.



Flowering Plants

Monkshood (*Aconitum columbiana*)

The distinctive flowers of the moisture loving monkshood are typically blue, but a white flowered form is fairly common. The foliage of monkshood is similar to the closely related western larkspur. All parts of the monkshood plant are extremely toxic if eaten. Poisoning symptoms include general weakness, and eventual respiratory paralysis. Abdominal pain and nausea may also occur. A fatal dose for an average dog is 0.2 ounces (5 grams) and horses are killed by doses as small as twelve ounces (350 grams).



Baneberry (*Actea rubra*)

Baneberry is found in wetlands and commonly grows to three feet tall. All parts of this plant are poisonous if eaten, but the roots and berries are especially toxic. While death from baneberry poisoning is rare, symptoms can be extremely uncomfortable and include vomiting, delirium, and stomach cramps lasting up to three hours. Symptoms may be worse for children than adults. The berries of the baneberry plant are bright and colorful and are especially attractive to children. Baneberry was used as a medicinal plant by American Indians.



Dogbane (*Apocynum androsaemifolium*)

Dogbane is a low-growing plant common in a wide range of dry habitats. It has milky sap and pink, waxy flowers. The toxic compound is a cardiac glycoside, a chemical that can cause heart failure. Dogbane is especially toxic to dogs. The bitter taste of this plant makes accidental poisoning unlikely. This plant is sometimes used in herbal remedies and its misuse can cause poisoning. A similar species called Indian hemp (*Apocynum cannabinum*) is tall, upright, and commonly found near low elevation wetlands. It too contains the cardiac glycoside found in dogbane. Indian hemp was used as a fiber source for rope and twine by American Indians.



Milkweed (*Asclepias speciosa*)

Milkweed is typically found in dry habitats, especially in dry, disturbed areas like roadsides, pastures, and dry streambeds. Although eating milkweed is rarely fatal to humans (because it tastes nasty), livestock occasionally die from eating it. Young shoots of milkweed can be eaten, but only after thorough cooking. Never eat parts of mature plants. Milkweed sap can cause a rash on people with sensitive skin. However, milkweed is the only thing the larvae of monarch butterflies eat. So, even though it is poisonous to humans & livestock it is a very important plant to the monarch butterfly.




Stinging Nettle (*Urtica dioica*)

Stinging nettle is common in wet areas throughout the forests of southeast Idaho. This thin, upright nettle usually occurs in dense patches. Stinging nettle causes extreme discomfort when touched. Tiny hairs on the stem and leaves break off readily and leave small amounts of a very irritating chemical in your skin. People differ greatly in their sensitivity to stinging nettle and different areas of your body may be more sensitive to it. Young stinging nettle leaves can be eaten if cooked properly. Never eat nettle flowers.



Corn Lilly (*Veratrum spp.*)

Corn lily is common in forested areas where the soil is consistently moist. It is a tall, erect plant, all parts of which are highly toxic and potentially fatal. The highest concentration of toxins are in the roots. Early spring foliage seems to be more poisonous than mature leaves. Poisoning symptoms include burning sensations in the mouth, vomiting, diarrhea, sweating, blurred vision, hallucinations, and general paralysis. It may cause birth defects if consumed by pregnant females. The occasional cases of human poisoning have been attributed to misuse of medicinal preparations. *Veratrum* species are used medicinally to treat high blood pressure though doses are difficult to standardize. This plant was used medicinally by American Indians for external afflictions and used in cleansing rites as a purgative.



Night shade (*Solanum dulcamara*)

The deadly nightshade is a common, slender vine often found in low elevation streambeds and other shaded, wet places. The entire plant, especially its unripe fruit, is toxic and potentially fatal. Nightshade most often poisons children who have been tempted to eat the bright-red, ripe berries. Although the unripe, green fruit is more toxic than the ripe fruit, children have died from consuming both (so don't take any chances). The toxicity of nightshade varies with the growing conditions of plant. Symptoms of poisoning include abdominal pain, thirst, restlessness, flushing, and skin irritation. In severe cases, victims may experience vomiting, difficulty breathing, dilated pupils, diarrhea, bloody urine, loss of sensation, and even death.




Locoweeds and Milkvetches (*Astragalus* spp.)

The genus *Astragalus* is the most diverse group of flowering plants in the world. Many species of *Astragalus* are valuable forage for livestock (milkvetches), but a few varieties (the locoweeds) are poisonous. Locoweeds pose little danger to humans as the toxic effects generally requires consuming large quantities of plant material (the seeds and pods are more toxic). Locoweeds can be addictive to horses. Animals "hooked" on locoweed will commonly display noticeable changes in behavior, may shun all other foods, and may eventually die.



Tansy (*Tanacetum vulgare*)

Tansy (not to be confused with tansy ragwort) is a widespread, tall, strong scented herb, with deeply toothed leaves. It is common in disturbed areas at low elevations. The entire plant is toxic and capable of causing severe illness and sometimes death. Tansy is sometimes misused as an herbal tea. Poisoning symptoms include rapid and feeble pulse, stomach lining inflammation, spasms and convulsions. The dried leaves and flowers have been used to kill intestinal worms, promote menstruation, and cause abortion.



Water Hemlock (*Cicuta douglasii*)

Common in damp ground and shallow water, the water hemlock can be identified by the veins in the leaves that terminate at the bottom of the serrations instead of the end of the serrations as in most plants. This is the most poisonous plant in North America; unfortunately, it is sometimes mistaken for look-alike edibles. The entire plant is fatally poisonous if ingested but the root is most toxic (a piece of root the width of a finger can kill an adult human). When cut, the root exudes a yellow juice that contains the toxin. Symptoms occur within 15 minutes to one hour after ingestion and include nausea, excessive salivation and frothing at the mouth, vomiting, violent convulsions, and usually death.



Poison hemlock (*Conium maculatum*)

Introduced from Europe, poison hemlock is easily identified by its purple spotted stems. The plant emits a mousy smell when bruised or crushed. Poison hemlock grows in disturbed soils and is often weedy. The entire plant is highly toxic, especially for children. It is more toxic in warm, sunny weather, and young foliage is more toxic than mature foliage. Symptoms of poisoning include an initial stimulation of the nervous system followed by severe depression of nervous system responses, slowing of the heart, paralysis, and eventually respiratory paralysis. Small quantities of poison hemlock ingested by pregnant females may cause birth defects. This is the plant used to silence the philosopher Socrates in 339 BC.



Elderberry (*Sambucus spp.*)

Elderberries can be found throughout southeastern Idaho. The leaves, stems, bark, and roots contain compounds that are strongly purgative and cyanide producing. Eating uncooked berries may cause nausea, vomiting, and diarrhea. Children can get sick from using hollow elder stems for peashooters or whistles. Elderberry flowers and fruits are edible when cooked and are commonly used to make wine and jelly. Elderberry stems and roots were used medicinally as an emetic by American Indians.



Hound's Tongue (*Cynoglossum officinale*)

Hound's Tongue is an attractive plant with dull reddish or purple flowers. It is common in pine and Douglas-fir forests. Eating any part of the hound's tongue can cause permanent and irreversible liver damage.



Tansy Ragwort (*Senecio jacobaea*)

All parts of this yellow-flowered, wasteland weed are toxic, potentially carcinogenic, and may cause severe liver damage. This plant is seldom immediately fatal to humans. Milk from cows that have eaten tansy ragwort and honey from bees that pollinate its flowers can cause the same ill effects as consuming the plant directly. Livestock that eat tansy ragwort over extended periods may die.



Buttercups (*Ranunculus spp.*)

There are many species of buttercups in our area. They typically have shiny yellow flowers and are most often found in areas with wet or damp soils. Fresh plants often contain irritant oils which can cause blistering of the mouth and digestive tract though this is almost never fatal. Some buttercup species are more toxic than others and some species are used as medicinal herbs. Buttercups are considered more toxic to grazing animals than humans.



Larkspur (*Delphinium spp.*)

There are many species of larkspur in our area. The most common ones are blue, but white flowered forms are not uncommon. A tall species (western larkspur) is found along streams in forested areas. A short species, (Nuttall's larkspur) is found in dry soils in both forests and shrub lands. Symptoms of poisoning include general weakness and eventually respiratory paralysis. Abdominal pain and nausea are also commonly experienced. Young plants and seeds contain the highest amounts of toxins and poisonings are more common in animals than humans.



Poison Ivy (*Rhus radicans*)

Poison ivy, our most familiar poisonous plant, occurs in many low and mid-elevation habitats. Its growth form can vary from a low shrub to a traveling vine. The entire plant (stems, flowers, pollen, and even the smoke of burning plants) is allergenic, causing severe and continuing skin irritation on contact for most people. You can also be poisoned by secondary contact with tools or clothes that have touched poison ivy. The fluid in poison ivy induced blisters contain the compound that causes the infamous reaction and can cause further contamination (so don't pop those blisters).



Steer's Head (*Dicentra uniflora*)

Related to the gardener's bleeding hearts, steer's head grows in rich, well-drained, organic soils throughout the northwest. Death in humans is rare but symptoms may include vomiting, diarrhea, severe trembling, and difficulty in breathing. Children are at greater risk than adults.



Spurge (*Euphorbia spp.*)

There are many species of spurge. In our area, the noxious leafy spurge (*E. esula*) is most common. Plants bleed a milky juice when cut or bruised. People have died from consuming dried plants, fruits, and seeds. Symptoms of spurge poisoning include convulsions, burning at the mouth, fluid build-up in lungs, and constriction of pupils. Even licking your fingers after handling the plants can cause burning lips and tongue. The milky sap of spurge may cause a rash or blistering in people with sensitive skin.



Klamath Weed or St. Johnswort (*Hypericum perforatum*)

Klamath weed is an erect herb with opposite leaves, and bright yellow flowers found in dry areas at low elevations. This has minute oily droplets on the stems that can cause a skin rash on people with sensitive skin. Consuming large quantities of Klamath weed may cause convulsions, increased heart-rates and temperature, diarrhea, and even blindness in both humans and animals. Animals often survive Klamath weed poisoning only to die from secondary complications, such as failure to eat. This plant has long been used in herbal remedies, and was recently identified as an effective treatment against some forms of depression.



Lupines (*Lupinus spp.*)

Lupines are found in a wide range of habitats. All lupines share the characteristic leaf shape seen in the accompanying picture. All parts of lupines are toxic, especially spring foliage, flowers, and fruits. Lupine consumption by pregnant females can cause birth defects. Documented cases show cows and goats that eat lupines can pass toxins through their milk. Deformities in puppies and human babies have been linked to lupine contaminated milk consumed by pregnant females. Death from lupine poisoning is rare among humans but livestock that eat lupine in great quantities may die.



Chokecherry (*Prunus virginiana*)

Chokecherry is found throughout southeast Idaho. Although the fruits of cherries (Chokecherry and other types of *Prunus* species) are enjoyed by literally thousands of people every year, there have been a few cases of children being poisoned by eating large number of seeds along with the cherries. Cherry seeds, leaves, bark, and shoots contain cyanide producing compounds. These compounds have caused livestock deaths. Cyanide poisoning initially causes rapid breathing followed by slow and difficult breathing, anxiety, confusion, headache, low blood pressure and rapid heart rate. Convulsions, coma, and death can occur rapidly. Chokecherries are popular in this area for making jams and syrups.



Chokecherry